

RESEARCH

Open Access



Up-and-down determination of prophylactic norepinephrine boluses combined with crystalloid co-load for preventing post-spinal anesthesia hypotension during cesarean section

Chengfei Xu^{1†}, Peipei Zhang^{2†}, Chunying Dai¹, Jingjun Zhang¹, Honghao Wu¹, Qianying Liu¹, Zhiqiang Zhang¹, Fengwei Yin¹, Huadong Ni^{3*} and Liang Cheng^{1*}

Abstract

Background The use of a fluid co-load has been shown to enhance hemodynamic stability and diminish the occurrence of hypotension after spinal anesthesia when paired with prophylactic norepinephrine. This research aimed to identify the effective dosages (ED₉₀ and ED₅₀) of prophylactic norepinephrine boluses, in conjunction with a crystalloid co-load, for the prevention of hypotension after spinal anesthesia in cesarean delivery patients.

Methods Patients were administered crystalloid co-loads at a dosage of 10 mL/kg, in addition to preventive norepinephrine dosages direct following spinal anesthesia administration. The dosages of norepinephrine were established employing the up-and-down sequential allocation technique, starting with 8 µg and progressively rising by 1 µg increments. The primary objective was to detect the effective dosage (ED₉₀ and ED₅₀) of norepinephrine necessary to avoid hypotension following spinal anesthesia.

Results The ED₉₀ for a single norepinephrine bolus, in combination with a crystalloid co-load, was calculated to be 5.35 µg (95% CI: 4.75 to 7.13). The ED₅₀ was determined to be 4.05 µg (95% CI: 3.68 to 4.46) using the up-and-down method and 3.926 µg (95% CI: 3.362 to 4.422) through the probit regression model.

Conclusion A prophylactic norepinephrine bolus of 5.35 µg, administered with a crystalloid co-load, effectively prevents hypotension following the spinal anesthesia in cesarean delivery patients.

Keywords Crystalloid co-load, Norepinephrine, Post-spinal anesthesia hypotension, Cesarean section

[†]Chengfei Xu and Peipei Zhang contributed equally to this work.

*Correspondence:
Huadong Ni
huadongni@zjxu.edu.cn
Liang Cheng
bbchengliang@163.com

¹Department of Anesthesiology, The Third People's Hospital of Bengbu, 38 Shengli Middle Road, Bengbu 233000, China

²Department of Obstetrics and Gynecology, The Third People's Hospital of Bengbu, 38 Shengli Middle Road, Bengbu 233000, China

³Department of Anesthesiology and Pain Research center, The Affiliated Hospital of Jiaxing University, 1882 Zhonghuan South Road, Jiaxing 314001, China



Introduction

Spinal anesthesia is the favored technique for cesarean delivery, yet it is often accompanied by hypotension after the spinal anesthesia, which remains the most prevalent complication related to maternity. This hypotension can lead to serious issues such as nausea, vomiting, reduced placental perfusion, and fetal acidosis [1]. To both prevent and manage this condition, vasopressors and fluid loading are essential, as these measures help stabilize hemodynamics [2, 3]. Fluid loading is particularly vital in addressing post-spinal anesthesia hypotension by maintaining intravascular volume, promoting sufficient tissue perfusion, and mitigating the hypotensive effects of sympathetic blockade [4]. Furthermore, fluid loading lessens dependence on vasopressors while enhancing the hemodynamic stability achieved through prophylactic vasopressor administration [3, 5].

The timing of fluid loading is critical for maximizing its effectiveness. Historically, preloading—conducted 15 to 30 min before spinal anesthesia—was commonly used during cesarean Sect. [6]. However, research has indicated that a 30 mL/kg crystalloid preload is inadequate in preventing hypotension following spinal anesthesia [7]. Consequently, clinical practice has increasingly favored co-loading, where fluids are administered immediately after spinal anesthesia induction. Co-loading has shown greater efficacy in augmenting intravascular volume during the time of maximal arterial expansion resulting from sympathetic blockade [8]. Recent studies have confirmed that co-loading with 10 mL/kg of crystalloids, together with prophylactic vasopressor infusion, successfully prevents post-anesthetic hypotension [9]. Nonetheless, the precise effective doses (ED_{90} and ED_{50}) of prophylactic norepinephrine in conjunction with a crystalloid co-load remain uncertain [10, 11].

Despite guidelines supporting prophylactic infusions to avert spinal hypotension [3], many anesthesiologists continue to prefer intravenous bolus administration [12]. Additionally, intermittent intravenous norepinephrine boluses have proven to be an effective strategy for avoiding hypotension caused by spinal anesthesia in obstetric patients, with minimal adverse effects [13]. In this investigation, we applied the up-and-down sequential allocation technique to ascertain the ED_{90} and ED_{50} of prophylactic norepinephrine boluses, administered alongside crystalloid co-loading, to prevent hypotension after the spinal anesthesia in the course of cesarean delivery.

Patients and methods

This prospective, double-blind, sequential allocation dose-finding study was authorized by the Research Ethics Board of the Third People's Hospital of Bengbu, China (Approval No. 2024-k21; June 24, 2024). Conducted between July 2024 and August 2024, the study utilized an

up-and-down design to determine dose efficacy [14]. The study enrolled non-laboring women at term scheduled for elective cesarean delivery. Written informed permission was obtained from everyone involved in the study, and the trial had been registered in ClinicalTrials.gov (NCT06498115, 2024/07/11).

Study population

Inclusion criteria for participants included a healthy singleton pregnancy at full term (above 37 weeks of pregnancy), elective cesarean delivery under spinal anesthesia, the weight of the participant ranging from 50 to 100 kg, American Society of Anesthesiologists (ASA) physical status classification of I or II [15], height ranging from 150 to 180 cm, and a fasting duration exceeding 6 h. To ensure adherence to current standardized guidelines for patient evaluation, the most recent European Society of Anaesthesiology and Intensive Care (ESAIC) Guidelines were also referenced [16]. Exclusion criteria included allergies or hypersensitivity to norepinephrine, hypertension, cardiovascular diseases, arrhythmias, pre-eclampsia, fetal abnormalities, spinal cord malformations, and patient refusal to participate.

Monitoring and anesthesia

On the morning of the surgery, baseline systolic and diastolic blood pressure (SBP and DBP) and heart rate (HR) were detected in the supine position using non-invasive methods. Initial SBP was established as the average of three successive readings obtained at 1-minute intervals employing an automatic blood pressure monitor. Blood pressure was measured at baseline and then at 1-minute intervals during the first 10 min after spinal anesthesia to monitor rapid hemodynamic changes. Thereafter, measurements were taken every 5 min until the end of the surgical procedure. During the procedure, continuous standard monitoring was performed, pulse oximetry, like non-invasive blood pressure, and electrocardiography using a monitoring system for the patient (MX550; Philips, Netherlands). Premedication was not administered. Participants were positioned in the left lateral decubitus position before spinal anesthesia. A 16-gauge epidural needle was used to puncture the L3-L4 interspace, followed by the insertion of a 25-gauge Whitacre needle. Upon confirmation of cerebrospinal fluid outflow, indicating successful subarachnoid space entry, 0.6% ropivacaine was intrathecally injected at a rate of 0.1 mL/sec. The ropivacaine solution was prepared by diluting 1.5 mL of 1% ropivacaine (Shijiazhuang Four Drugs Co., Ltd.) with 1 mL of cerebrospinal fluid, a dosage based on previous studies [17–19]. After the injection, patients were situated supine with a 15-degree left lateral tilt to achieve left uterine displacement [20]. Oxygen was administered at 5 L per minute via mask. The degree of the spinal

sensory blockage was evaluated employing the pinprick technique and verified to fall within the interval of thoracic (T) 4–6. Investigation eliminated individuals who failed to meet this criterion.

Study protocol

A crystalloid preload was supplied at a dosage of 5 ml/kg over 20 min after an 18-gauge intravenous catheter was inserted into the right forearm prior to anesthesia administration. After spinal anesthesia was initiated, a crystalloid co-load solution consisting of complex sodium chloride (0.85% NaCl, 0.03% KCl, and 0.033% CaCl₂) was infused at 10 ml/kg over 10–15 min, then a preservation infusion rate of 0.1 ml/kg/min was conducted. To produce the investigation bolus dosage, the anesthesiologist assistant, who did not participate in patient treatment or gathering information but was responsible for dose preparation, added 2 mg of norepinephrine (Yuanda Pharmaceutical Co., Ltd.) to 500 ml of physiological saline. There was blinding of the patient and the participating anesthesiologist to group assignment, and the anesthesiologist in charge of collecting data was likewise blinded to the specific medication type and dose.

Based on prior research [21, 22], the first participant underwent a primary prophylactic bolus dose of 8 µg norepinephrine simultaneously with spinal anesthesia. This approach, validated in a randomized, double-blinded study comparing norepinephrine and phenylephrine, highlights the ability of an 8 µg norepinephrine bolus to maintain cardiac output and reduce the incidence of bradycardia [23]. In our study, this dosage was adopted to build upon prior evidence and ensure optimal maternal hemodynamic stability during elective cesarean sections. Subsequent norepinephrine doses were adjusted in 1 µg increments, depending on their effectiveness in avoiding hypotension after spinal anesthesia, defined as an SBP drop below 80% of baseline. The up-and-down sequential allocation approach directed these modifications [24, 25]. This method is a sequential allocation technique designed to estimate the ED₅₀ and ED₉₀ based on individual patient responses to varying doses of norepinephrine. The randomization process in our study was based on the patient's response to the initial dose. Initially, the patient was administered a starting dose of 8 µg of norepinephrine. If the patient experienced hypotension, the dose was adjusted increased by 1 µg for the next patient. If the patient did not experience hypotension, the dose was downward by 1 µg for the next patient. This “Up-and-Down” process continued until sufficient data was collected to estimate the effective dose range. This method allows for rapid adjustments in response to individual variability, enabling us to identify the dose range that most effectively prevents hypotension during spinal anesthesia for cesarean section.

Outcome measurement

The primary finding was to determine the effective doses of prophylactic norepinephrine essential to avoid hypotension following the spinal anesthesia in 90% (ED₉₀) and 50% (ED₅₀) of patients. Hypotension was known as a SBP falling below 80% of baseline, while the goal was to maintain SBP within 95% of baseline. An increase in SBP of 20% or more from baseline was considered hypertension. Successful treatment was identified as achieving an SBP within 95% of baseline within 1 min post-administration. Provided that the SBP kept below 80% of the initial level, 6 mg of ephedrine was provided. Treatment for bradycardia, which is defined as a pulse rate under 50 beats per minute, included the administration of 0.5 mg of atropine. Systematic documentation of maternal consequences, like infusion volume, bleeding, urination, and adverse reactions (like bradycardia, nausea, vomiting, hypertension, and hypotension), was conducted. Furthermore, the study assessed the fetal heart rates prior to and following spinal anesthesia, the Apgar ratings at 1 and 5 min, and the umbilical arterial blood gases.

Statistical analysis

Due to the non-independent and unknown data distribution inherent in the up-and-down study design, theoretical guidelines for precise sample size calculations to estimate the effective dose interval ED₉₀ were not applicable [26]. Simulation studies indicate that a sample size of 20–40 patients generally yields stable dose estimates across various scenarios [26]. To enhance the reliability of our results, we included 51 patients in the study. The ED₅₀ was determined with a 95% confidence interval (CI) using the up-and-down method. Probit regression analysis was also conducted to estimate both ED₅₀ and ED₉₀. Visualization of data was conducted employing version 9.5.1 of GraphPad Prism, and statistical analysis was done utilizing version 20.0 of the SPSS program.

Results

Patient recruitment

Figure 1 provides a detailed flow chart outlining the enrollment, exclusion, follow-up, and analysis of participants. Following the application of exclusion criteria and completion of follow-up techniques, 51 out of the 60 patients originally planned for optional cesarean sections using spinal anesthesia fulfilled the last criteria for inclusion and were enrolled in the investigation. Table 1 summarizes maternal demographics and surgical characteristics. Participants average age was 30.55 ± 5.07 years, and mean body mass index (BMI) was 29.12 ± 3.36 kg/m². Baseline SBP and DBP were 125.51 ± 6.84 mmHg and 74.65 ± 4.45 mmHg, respectively. The spinal block level ranged from T4 to T6, and the average duration of surgery was 49.20 ± 6.47 min.




Fig. 1 Flowchart depicting the patient recruitment process, including stages of enrollment, exclusion criteria, follow-up procedures, and final analysis

Table 1 Patient demographics and surgical characteristics

Characteristics	Index (n=51)
Age (years)	30.55 ± 5.07
Weight (kg)	76.96 ± 8.37
Height (cm)	162.70 ± 4.85
Body mass index (kg/m ²)	29.12 ± 3.36
Gestation (weeks)	38.21 ± 0.99
SBP at baseline (mmHg)	125.51 ± 6.84
DBP at baseline (mmHg)	74.65 ± 4.45
HR at baseline (beats/min)	84.16 ± 12.57
Block level (T)	T5 (T4, T6)
Induction to delivery (min)	11.20 ± 2.50
Duration of surgery (min)	49.20 ± 6.47

SBP, systolic blood pressure; DBP, diastolic blood pressure; HR, heart rate. Data are presented as mean ± SD and median (IQR)

Primary outcomes

Figure 2 illustrates the sequences of patient responses to prophylactic norepinephrine dosages, categorized as effective or ineffective based on the up-and-down sequential technique. Figure 3 presents dose-response curves for prophylactic norepinephrine, which have been obtained using probit regression analysis. Isotonic regression analysis determined the ED₅₀ for prophylactic norepinephrine to be 4.05 µg (95% CI: 3.68 to 4.46). Probit regression analysis estimated the ED₅₀ to be 3.926 µg (95% CI: 3.362 to 4.422) and the ED₉₀ to be 5.35 µg (95% CI: 4.75 to 7.13). Table 2 details the response rates for various prophylactic norepinephrine doses.

Maternal and neonatal outcomes

Patients received a crystalloid pre-load of 365 ml and a crystalloid co-load of 730 ml, as detailed in Table 3. Maternal outcomes are summarized in Table 4. Hypotension occurred in 23 patients (45.10%), while 1 patient experienced hypertension. Additionally, 3 patients reported nausea, 2 experienced vomiting, and 3 exhibited bradycardia. Neonatal outcomes are detailed in Table 5. A range of 3.14 ± 1.88 for umbilical artery base excess values and 7.33 ± 0.04 for pH values were detected. All pH values were within the normal range [27], with no neonates presenting with fetal acidosis (pH < 7.2). At both 1 min and 5 min, all newborns achieved Apgar scores of 8 or above.

Discussion

Our study determined the effective dose of norepinephrine required to prevent hypotension following the spinal anesthesia, establishing the ED₉₀ as 5.35 µg (95% CI: 4.75 to 7.13) when administered with a crystalloid co-load. This finding highlights a significant reduction in norepinephrine dose compared to previous studies [13]. The ED₅₀ in our study was approximately 4 µg, as estimated by isotonic regression analysis (4.05 µg, 95% CI: 3.68 to 4.46) and probit regression analysis (3.926 µg, 95% CI: 3.362 to 4.422), which is notably lower than ED₅₀ values reported in earlier research [28].

Fluid loading, whether used independently or in combination with vasopressors, is a well-established approach for avoiding and managing post-spinal anesthesia




Fig. 2 Graph showing the distribution of patient responses to different norepinephrine bolus doses, categorizing them as either effective or ineffective




Fig. 3 Dose-response curve for norepinephrine: This graph illustrates the relationship between various norepinephrine dosages and the proportion of participants achieving effective treatment, identified as the restoration of systolic blood pressure (SBP) to within 95% of baseline values. The ED_{50} and ED_{90} values were calculated employing probit regression analysis

Table 2 Response rates for doses of prophylactic norepinephrine

Assigned dose of norepinephrine (ug)	Number of successes	Total number	Response rate (%)
2	0	3	0.00
3	3	11	27.27
4	9	19	47.37
5	11	13	84.62
6	3	3	100.00
7	2	2	100.00
8	1	1	100.00

Data are presented as number (%)

Table 3 Volumes of patient fluid intake and output

Parameters	Index (n = 51)
Crystalloid pre-load (ml)	365 (332.50, 410.00)
Crystalloid co-load (ml)	730 (665.00, 820.00)
Bleeding volume (ml)	246.08 ± 93.19
Urine volume (ml)	119.61 ± 51.07

Data are presented as mean ± SD and median (IQR)

Table 4 Maternal outcomes

Outcome	Index (n = 51)
Hypotension	23 (45.10%)
Hypertension	1 (1.96%)
Nausea	3 (5.88%)
Vomiting	2 (3.92%)
Bradycardia	3 (5.88%)

Data are presented as number (%)

Table 5 Neonatal umbilical artery outcomes

Outcome	Index (n = 51)
PO ₂ (mmHg)	21.39 ± 3.27
PCO ₂ (mmHg)	51.22 ± 6.14
pH	7.33 ± 0.04
HCO ₃ ⁻ (mmol/l)	22.65 ± 1.42
Base excess (mmol/l)	-3.14 ± 1.88
Apgar score at 1 min	9.00 (8.00, 9.00)
Apgar score < 7 at 1 min, n (%)	0 (0)
Apgar score at 5 min	10.00 (9.00, 10.00)
Apgar score < 7 at 5 min, n (%)	0 (0)

Data are presented as mean ± SD, median (IQR) and number (%)

hypotension after spinal [29]. Various fluid loading techniques, differing in their nature (crystalloid vs. colloid), time (preload vs. co-load), and volume, have demonstrated efficacy in reducing, but not entirely eradicating, post-spinal hypotension. The Vasopressor usage, therefore, plays a vital role in enhancing outcomes [30]. Chen et al. [31] have proposed that colloid co-loading does not provide any further advantages compared to crystalloid co-loading when used with prophylactic norepinephrine infusions. This supports the recommendation for

crystalloid in large co-load volumes, like 10 mL/kg, for optimal outcomes. Our study's use of a 10 mL/kg crystalloid co-load aligns with these findings. While a crystalloid preload can partially address pre-operative hypovolemia and improve cardiac output prior to spinal anesthesia, it often fails to fully maintain systolic blood pressure (SBP) and correct flow time, thus not entirely eliminating post-spinal hypotension [32, 33]. In contrast, a crystalloid co-load offers more efficient growth of volume inside the vessel with slower redistribution [34]. Colloids, owing to their greater molecular size, tend to remain in the intravascular space longer compared to crystalloids, enhancing intravascular volume growth and osmotic pressure [35, 36]. Recently, a study conducted by Theodoraki et al. revealed that the incidence of hypotension after spinal anesthesia was equally low whether utilizing either colloid preload or crystalloid co-load in conjunction with infusion of prophylactic norepinephrine [37]. This finding underscores the possible benefits of employing crystalloids in this particular scenario.

While prophylactic infusion of vasopressors is commonly recommended to manage spinal hypotension and stabilize hemodynamic fluctuations, it is connected with an elevated risk of reactive hypertension [3]. In contrast to bolus regimens, infusion procedures often need a greater total dosage of phenylephrine to sustain maternal arterial blood pressure at the initial level during the pre-delivery interval [38]. The bolus approach allows for the immediate administration of an efficient dosage of phenylephrine, which can rapidly restore maternal vascular resistance throughout spinal blockade. As a result, many anesthesiologists prefer bolus administration and are accustomed to providing frequent dosages rather than initiating an infusion throughout spinal anesthesia [39]. In this study, we employed a norepinephrine bolus, a method familiar to most anesthesiologists. Currently, there is limited research specifically defining the ED₅₀ and ED₉₀ of norepinephrine used with crystalloid co-loading for optional cesarean delivery using spinal anesthesia. Norepinephrine boluses application in conjunction with crystalloid co-loading for hypotension management in this context remains inadequately explored.

Previous studies have provided varying estimates for the ED₉₀ of prophylactic norepinephrine combined with crystalloid co-loading. ED₉₀ values for prophylactic norepinephrine boluses were determined by Onwochei et al. [13], to be 5.49 µg (95% CI: 5.15–5.83 µg) and 5.80 µg (95% CI: 5.01–6.59 µg). Their study utilized lactated Ringer's solution for crystalloid preload, introduced as a bolus infusion at 10 mL/kg with a maximum volume of 1000 mL via a pressure bag set at 250 mm Hg. It is important to note that their study focused on preload rather than co-load. Network meta-analyses have indicated that crystalloid co-loading is more effective than crystalloid

preload in managing spinal hypotension during elective cesarean Sect. [5]. Preload volumes can rapidly redistribute into the interstitial space due to sympathetic blockade, which may lead to atrial chamber distension and increased secretion of atrial natriuretic peptide [40]. This peptide causes peripheral vasodilation and diuresis. In contrast, co-loading can reduce these effects by minimizing fluid redistribution into the interstitial space and decreasing hydrostatic pressure during spinal-induced vasodilation [37, 41]. Guo et al. [22] discovered the ED₉₀ for a bolus of prophylactic norepinephrine, administered with hydroxyethyl starch (130/0.4) co-load, to be 8.0 µg (95% CI: 7.1 to 11.0 µg). This higher ED₉₀ might be attributable to the relatively smaller co-load volume of 500 mL used in their study. Their findings also suggested no significant difference between colloid and crystalloid co-loading [31].

This discussion integrates findings from our study with existing literature, highlighting the relevance of crystalloid co-loading and bolus administration in managing spinal hypotension during cesarean sections. Further research is needed to refine these parameters and optimize clinical practice. This investigation has some restrictions that should be considered. First, the investigation concentrated exclusively on the initial incidence of hypotension following spinal anesthesia and did not account for subsequent episodes throughout the surgical procedure, which may have distinct response patterns. Second, variability in individual sensitivity to vasoactive agents could have influenced the study's outcomes. Third, the geographic homogeneity of the study population may limit the generalizability of the results to more diverse and broader populations. Finally, the lack of a control group constrains our ability to establish definitive causal relationships.

This investigation has some restrictions that should be considered. First, the investigation concentrated exclusively on the initial incidence of hypotension following spinal anesthesia and did not account for subsequent episodes throughout the surgical procedure, which may have distinct response patterns. Second, the high incidence of hypotension (45.10%) observed in our study warrants further discussion. One contributing factor may be the pharmacodynamic properties of norepinephrine, which has an onset time of 1–2 min. In our study, a treatment was considered successful if systolic blood pressure (SBP) returned to ≥95% of baseline within 1 min after norepinephrine administration. This strict criterion may have categorized cases as hypotensive before the drug fully took effect. Furthermore, norepinephrine's short duration of action increases the likelihood of transient hypotensive episodes between doses, especially if redosing is delayed. This time-sensitive nature of norepinephrine administration reflects the challenges in managing

maternal hemodynamics during spinal anesthesia. These factors likely contributed to the high incidence of hypotension in our study, highlighting an inherent limitation of the study protocol and the practical challenges in optimizing maternal hemodynamics with vasopressor boluses. Third, variability in individual sensitivity to vasoactive agents could have influenced the study's outcomes. Finally, the geographic homogeneity of the study population may limit the generalizability of the results to more diverse and broader populations.

In conclusion, this study determines that the ED₉₀ of norepinephrine administered with a crystalloid co-load is 5.35 µg (95% CI: 4.75 to 7.13), based on probit regression analysis. The ED₅₀ values for a single norepinephrine bolus to avoid hypotension throughout elective cesarean section were found to be 4.05 µg (95% CI: 3.68 to 4.46) using the up-and-down method and 3.926 µg (95% CI: 3.362 to 4.422) according to the probit regression model.

Acknowledgements

Not applicable.

Author contributions

CX and PZ conducted the majority of the experiments and wrote the manuscript. CD, JZ and HW conceived and designed the study. QL, and ZZ performed the data analysis. FY collected the data. LC and HN coordinated and supervised the experiments. All authors read and approved the final manuscript.

Funding

The current study was supported, in part, by grants from Research Project on the Inheritance and Innovation of Traditional Chinese Medicine of Anhui Province (2024CCCX272), Key Speciality of Anhui Province (2022-AH-105), Scientific research project of Anhui Provincial Health Commission (AHWJ2023A30069), Bengbu Science and Technology Bureau project (20220115), Natural Science Key Program of Bengbu Medical University (2023byzd113), Clinical Key Specialty of Zhejiang Province Anesthesiology (2023ZJZK001).

Data availability

Access to the datasets utilized and/or analyzed in the present work may be obtained from the corresponding author upon an acceptable request.

Declarations

Ethics approval and consent to participate

The present study was approved by the Research Ethics Board of the Third People's Hospital of Bengbu (Anhui, China) and registered in the ClinicalTrials.gov (NCT06498115, 2024/07/11). All of the participants who participated in the study provided written informed consent at the time of enrolment.

Consent for publication

Not applicable.

Competing interests

The authors declare no competing interests.

Received: 3 September 2024 / Accepted: 20 January 2025

Published online: 25 January 2025

References

1. Tawfik MM, Tarbay AI, Elaidy AM, Awad KA, Ezz HM, Tolba MA. Combined Colloid Preload and Crystalloid Coload Versus Crystalloid Coload during

- spinal anesthesia for cesarean delivery: a Randomized Controlled Trial. *Anesth Analg.* 2019;128(2):304–12.
2. Practice Guidelines for Obstetric Anesthesia. An updated report by the American Society of Anesthesiologists Task Force on Obstetric Anesthesia and the Society for Obstetric Anesthesia and Perinatology. *Anesthesiology.* 2016;124(2):270–300.
3. Kinsella SM, Carvalho B, Dyer RA, Fernando R, McDonnell N, Mercier FJ et al. International consensus statement on the management of hypotension with vasopressors during caesarean section under spinal anaesthesia. *Anaesth* 2018;73(1):71–92.
4. Ayebele ET, Kwizera A, Mijumbi C, Kizito S, Roche AM. Ringer's Lactate Versus Normal saline in Urgent Cesarean Delivery in a Resource-Limited setting: a pragmatic clinical trial. *Anesth Analg.* 2017;125(2):533–9.
5. Rijs K, Mercier FJ, Lucas DN, Rossaint R, Klimek M, Heesen M. Fluid loading therapy to prevent spinal hypotension in women undergoing elective caesarean section: Network meta-analysis, trial sequential analysis and meta-regression. *Eur J Anaesthesiol.* 2020;37(12):1126–42.
6. Loubert C. Fluid and vasopressor management for cesarean delivery under spinal anesthesia: continuing professional development. *Can J Anaesth = J Canadien D'anesthésie.* 2012;59(6):604–19.
7. Park GE, Hauch MA, Curlin F, Datta S, Bader AM. The effects of varying volumes of crystalloid administration before cesarean delivery on maternal hemodynamics and colloid osmotic pressure. *Anesth Analg.* 1996;83(2):299–303.
8. Kaufner L, Karekla A, Henkelmann A, Welfe S, von Weizsäcker K, Hellmeyer L, von Heymann C. Crystalloid coload vs. colloid coload in elective caesarean section: postspinal hypotension and vasopressor consumption, a prospective, observational clinical trial. *J Anesth.* 2019;33(1):40–9.
9. Chen Y, Xiong X, Qin R, Guo L, Shi Y, Ni X. Up-and-down determination of different crystalloid coload volumes on the ED 90 of prophylactic norepinephrine infusion for preventing Postspinal Anesthesia Hypotension during Cesarean Section. *Drug Des Devel Ther.* 2024;18:2609–16.
10. Hassabelnaby YS, Hasanin AM, Adly N, Mostafa MMA, Refaat S, Fouad E, et al. Comparison of two norepinephrine rescue bolus for management of post-spinal hypotension during cesarean delivery: a randomized controlled trial. *BMC Anesthesiol.* 2020;20(1):84.
11. Chen Z, Zhou J, Wan L, Huang H. Norepinephrine versus phenylephrine infusion for preventing postspinal hypotension during cesarean section for twin pregnancy: a double-blinded randomized controlled clinical trial. *BMC Anesthesiol.* 2022;22(1):17.
12. Allen TK, Muir HA, George RB, Habib AS. A survey of the management of spinal-induced hypotension for scheduled cesarean delivery. *Int J Obstetric Anesth* 2009;18(4):356–61.
13. Onwoche DN, Ngan Kee WD, Fung L, Downey K, Ye XY, Carvalho JCA. Norepinephrine Intermittent Intravenous boluses to prevent hypotension during spinal anesthesia for cesarean delivery: a sequential allocation dose-finding study. *Anesth Analg.* 2017;125(1):212–8.
14. Oron AP, Souter MJ, Flournoy N. Understanding Research methods: Up-and-down designs for dose-finding. *Anesthesiology.* 2022;137(2):137–50.
15. Doyle DJ, Hendrix JM, Garmon EH. American Society of Anesthesiologists Classification. StatPearls. Edn. Treasure Island (FL) ineligible companies. Disclosure: Joseph Maxwell Hendrix declares no relevant financial relationships with ineligible companies. Disclosure: Emily Garmon declares no relevant financial relationships with ineligible companies.: StatPearls Publishing Copyright © 2024. StatPearls Publishing LLC. 2024.
16. Lamperti M, Romero CS, Guarracino F, Cammarota G, Vetrugno L, Tufegdizic B et al. Preoperative assessment of adults undergoing elective noncardiac surgery: updated guidelines from the European Society of Anaesthesiology and Intensive Care. *Eur J Anaesthesiol* 2025;42(1):1–35.
17. Khaw KS, Ngan Kee WD, Wong EL, Liu JY, Chung R. Spinal ropivacaine for cesarean section: a dose-finding study. *Anesthesiology.* 2001;95(6):1346–50.
18. Chen XZ, Chen H, Lou AF, Lü CC. Dose-response study of spinal hyperbaric ropivacaine for cesarean section. *J Zhejiang Univ Sci B.* 2006;7(12):992–7.
19. Tang Y, Yang M, Fu F, Huang X, Feng Y, Chen X. Comparison of the ED50 of intrathecal hyperbaric ropivacaine co-administered with or without intrathecal dexmedetomidine for cesarean section: a prospective, double-blinded, randomized dose-response trial using up-down sequential allocation method. *J Clin Anesth.* 2020;62:109725.
20. Vallejo MC, Attallah AF, Elzamzamy OM, Cifarelli DT, Phelps AL, Hobbs GR, Shapiro RE, Ranganathan P. An open-label randomized controlled clinical trial for comparison of continuous phenylephrine versus norepinephrine infusion in prevention of spinal hypotension during cesarean delivery. *Int J Obstet Anesth.* 2017;29:18–25.
21. Wang T, He Q, Zhang W, Zhu J, Ni H, Yang R, Liu Q, Xu L, Yao M. Determination of the ED(50) and ED(95) of intravenous bolus of norepinephrine for the treatment of hypotension during spinal anesthesia for cesarean delivery. *Exp Ther Med.* 2020;19(3):1763–70.
22. Guo L, Xu X, Qin R, Shi Y, Xue W, He L, Ma S, Chen Y. Prophylactic norepinephrine and Phenylephrine Boluses to prevent Postspinal Anesthesia Hypotension during Cesarean Section: a randomized sequential allocation dose-finding study. *Drug Des Devel Ther.* 2023;17:1547–55.
23. Wang X, Mao M, Zhang SS, Wang ZH, Xu SQ, Shen XF. Bolus norepinephrine and phenylephrine for maternal hypotension during elective cesarean section with spinal anesthesia: a randomized, double-blinded study. *Chin Med J* 2020; 133(5):509–16.
24. Dixon WJ. Staircase bioassay: the up-and-down method. *Neurosci Biobehav Rev.* 1991;15(1):47–50.
25. Stylianou M. N Flournoy 2002 Dose finding using the biased coin up-and-down design and isotonic regression. *Biometrics* 2002;58(1):171–7.
26. Pace NL, Stylianou MP. Advances in and limitations of up-and-down methodology: a précis of clinical use, study design, and dose estimation in anesthesia research. *Anesthesiology.* 2007;107(1):144–52.
27. Filippi L, Pascarella F, Pini A, Cammalleri M, Bagnoli P, Morganti R et al. Fetal oxygenation from the 23rd to the 36th week of Gestation evaluated through the Umbilical Cord Blood Gas Analysis. *Int J Mol Sci* 2023;24(15).
28. Qian J, Zhao YP, Deng JL, Wang LZ, Xiao F, Shen B, Yao HQ. Determination of the relative potency of Norepinephrine and Phenylephrine given as infusions for preventing hypotension during combined spinal-epidural anesthesia for cesarean delivery: a randomized Up-And-down sequential allocation study. *Front Pharmacol.* 2022;13:942005.
29. Mercier FJ. Cesarean delivery fluid management. *Curr Opin Anaesthesiol.* 2012;25(3):286–91.
30. Sakata K, Yoshimura N, Tanabe K, Kito K, Nagase K, Iida H. Prediction of hypotension during spinal anesthesia for elective cesarean section by altered heart rate variability induced by postural change. *Int J Obstet Anesth.* 2017;29:34–8.
31. Chen Y, Xu X, Qin R, Guo L, Ni X. Comparison of crystalloid and colloid coload combined with norepinephrine prophylaxis on post-spinal anesthesia hypotension during cesarean delivery: a randomized sequential allocation dose-finding study. *Front Med.* 2023;10:1214598.
32. Tamilselvan P, Fernando R, Bray J, Sodhi M, Columb M. The effects of crystalloid and colloid preload on cardiac output in the parturient undergoing planned cesarean delivery under spinal anesthesia: a randomized trial. *Anesth Analg.* 2009;109(6):1916–21.
33. Lee A, Ngan Kee W. Effects of Vasoactive Medications and maternal positioning during cesarean delivery on maternal hemodynamics and neonatal acid-base status. *Clin Perinatol.* 2019;46(4):765–83.
34. Gunusen I, Karaman S, Ertugrul V, Firat V. Effects of fluid preload (crystalloid or colloid) compared with crystalloid co-load plus ephedrine infusion on hypotension and neonatal outcome during spinal anaesthesia for caesarean delivery. *Anaesth Intensive Care.* 2010;38(4):647–53.
35. Lewis SR, Pritchard MW, Evans DJ, Butler AR, Alderson P, Smith AF, Roberts I. Colloids versus crystalloids for fluid resuscitation in critically ill people. *Cochrane Database Syst Rev.* 2018;8(8):Cd000567.
36. Gousheh MR, Akhondzade R, Asl Aghahoseini H, Olapour A, Rashidi M. The effects of Pre-spinal Anesthesia Administration of Crystalloid and Colloid Solutions on hypotension in Elective Cesarean Section. *Anesth Pain Med.* 2018;8(4):e69446.
37. Theodoraki K, Hadzilia S, Valsamidis D, Kalopita K, Stamatakis E. Colloid Preload versus Crystalloid Co-load in the setting of Norepinephrine infusion during Cesarean Section: time and type of administered fluids do not Matter. *J Clin Med* 2023;12(4).
38. Doherty A, Ohashi Y, Downey K, Carvalho JC. Phenylephrine infusion versus bolus regimens during cesarean delivery under spinal anesthesia: a double-blind randomized clinical trial to assess hemodynamic changes. *Anesth Analg.* 2012;115(6):1343–50.
39. Langesaeter E, Rosseland LA, Stubhaug A. Continuous invasive blood pressure and cardiac output monitoring during cesarean delivery: a randomized, double-blind comparison of low-dose versus high-dose spinal anesthesia with intravenous phenylephrine or placebo infusion. *Anesthesiology.* 2008;109(5):856–63.

40. Lee JE, George RB, Habib AS. Spinal-induced hypotension: incidence, mechanisms, prophylaxis, and management: summarizing 20 years of research. *Best Pract Res Clin Anaesthesiol* 2017;31(1):57–68.
41. Jin WD, Mao JQ, Liu J, Liang G, Jiang C, Sheng ZM. Comparative dose-response study on the infusion of Norepinephrine Combined with Crystalloid Coload versus Colloid Coload for preventing hypotension during spinal anesthesia for Cesarean Delivery. *Drug design, development and therapy* 2022;16:2617–26.

Publisher's note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.